

Causes and Management of Hoarseness of Voice in Benign Laryngeal Disorders in Azadi Teaching Hospital

Dr. Ayad Mustafa Bahaalddin, FICMS-ENT

Dr. Ali Hussein Jamal, FICMS-ENT

Dr. Maroof Saki Baki, FICMS-ENT

Department of otolaryngology, Azadi teaching hospital

Abstract:

Background: Hoarseness remain one of the major complain of patients visiting Otolaryngology clinic and regarded as one of the intractable conditions for treatment. It is almost a benign condition and could be treated by medical or surgical treatment according to the severity of symptoms and duration of patient complain.

Patients and Methods: A prospective study had been carried out on 120 patients who attended ENT out patient clinic in Azadi teaching hospital from January 2018 to January 2019. All the cases had been diagnosed with either indirect laryngeal mirror or by flexible laryngoscope. The cases treated in the out patient as a conservative measures or the patients had been admitted and treated with a surgical maneuvers.

Results: The patients consisted of 55% (64) males and 45% (56) females with an average age of 40 years. In this study the incidence of chronic hoarseness (65.84%) was more common than acute hoarseness (34.16%) and acute laryngitis (26.66%) was the most common cause of acute hoarseness, while Vocal cord polyp was the most common cause of chronic hoarseness (22.5%).

Conclusions: The commonest cause of hoarseness was found to be acute laryngitis of Larynx which predominantly affects young males. The peak age incidence is between 21-30 years, Males are affected more than females

Keywords: dysphonia, hoarseness, acute and chronic laryngitis, vocal cord polyp and nodule, psychogenic dysphonia, Reinke's oedema

Introduction: . Hoarseness is a general term defined as altered rough voice which interfere with vibration ,approximation or movements of the vocal cords. Dysphonia is defined as an abnormal voice quality or difficulty in producing voice particularly in patients who are singers, are changes in pitch and abnormal pitch range. human larynx is a specialized organ that enables us to speak & communicate and modulate sound The main functions in protection of the upper airway, in deglutition, respiration and phonation Voice is the primary means of communication for humans both socially and in the workplace.. Voice production can

be thought of in terms of three components: generation of airflow, vocal cord vibration that produces sound and shaping of the sound to produce various resonances. The lips, tongue, teeth and palate provide the distinct sounds of speech. The lungs first supply adequate airflow to overcome the resistance of the adducted vocal cords. The vocal cords are finely tuned neuromuscular units that adjust pitch and tone by altering their position, length, tension and mass.(3). Sound production occurs as a result of the vibration of the mucosa at the medial edge of each vocal cord. Thus, any structural, inflammatory or neoplastic lesion of the vocal cord affects voice production and quality. _

PATIENTS AND METHODS

One hundred and twenty patients complaining from hoarseness were evaluated prospectively at the department of otolaryngology, at Azadi Teaching Hospital, Kirkuk - Iraq, from the period January 2018 to January 2019. The patients aged 3 years to 70 years, they were 66 males and 54 females The plan of

evaluation included the following steps: 1. History : A- * Name * Age * Sex * occupation *Residence *Telephone No. *Date of examination _B-Chief complain and duration: C- History- History of present illness ; including : duration and time course of hoarseness

Initiating factors e.g. URTI,
Trauma, psychological
upset...etc. Exacerbating and
relieving factors. D-Systemic
review including all the systems:

1-Respiratory 2.Cardiovascular
3.Gastroenterology.4Haematology
5-Neurology 6.Endocrinology

.E- Past medical history ;
pulmonary, cardiovascular
gastrointestinal ,neurological,
autoimmune, endocrine and
psychological disorders.

D- Past surgical history ; regarding
history of intubation, or any neck,
heart or pulmonary surgery. E-

Drug history F-History of trauma
Blunt or penetrating trauma to neck
G- Social history ; concerning
Cigarette smoking, alcohol intake
and feeding habits.

H-Family history of the same
condition

2- Physical examination*

Physical examination ; include

A-General examination for

signs of systemic disease
e.g. general condition of the
patient ,level of consciousness
and nutritional state B-Systemic
Examination including:

1-Respiratory.

2.Cardiovascular.

3-Abdominal .

4-Neurology and other systems
should also be examined carefully
including full otolaryngology
examination and laryngeal
examination starting from surface
and external examination and lastly
laryngeal internal examination by
both indirect laryngoscopy and
flexible endoscopic examination of
the larynx and the patient should be
carefully prepared for the
examination and the larynx should
be carefully anaesthetized in order
to prevent gag reflex and to be
properly examined and in order to
not miss any area of the larynx .

Visualization of larynx by indirect Laryngoscopy, and those who fail mirror examination (even with the application of topical anesthetic) are assessed by flexible nasolaryngoscopy. The patients were divided into 2 groups according to the duration of hoarseness; • Acute hoarseness (less than 2 weeks); Patients are examined by indirect laryngoscopy and if it was difficult due to strong gag reflex or difficult anatomy then the patient was examined by flexible laryngoscopy. If only a self-limiting lesion was seen e.g. Acute laryngitis or voice strain and there is no feature of serious pathology ; then no further investigation done. But if a mass or impaired mobility of vocal cords were detected then a plan of investigations was followed as in chronic hoarseness. • Chronichoarseness (more than 2 weeks); Patients are also assessed by indirect laryngoscopy and if it was difficult then by flexible laryngoscopy and general impression of the lesion was obtained. Direct laryngoscopy under general anesthesia should be done in all cases of chronic

hoarseness' (either for diagnostic or therapeutic purposes). The plan of investigations include: □ Baseline investigations: CBP , ESR, prothrombin time, partial thromboplastin time, lateral X-ray of neck and chest X-ray. Additional investigations ; includes: fasting blood sugar , thyroid function test, rheumatoid factor, barium swallow, sputum for AFB. These investigations are only done when there are associated features suggestive of other diseases like TB ; and goitre as a cause of hoarseness. In the present study, 74 patients were examined satisfactorily by indirect laryngoscopy. The remaining 46 patients who failed examination by indirect laryngoscopy even after application of topical anesthesia, were examined by flexible nasolaryngoscope, But this was found to be difficult in 7 patients due to either patient uncooperation or difficult anatomy. Direct laryngoscopy was required for 58 patients, either for diagnostic purposes (in 18 patients) or for therapeutic purposes (in 40 patients).

RESULTS

1 - Distribution of patients according to duration of hoarseness: The patients were divided according to the duration of hoarseness into acute

were more in compares with cases with acute hoarseness(34.16).

2- Causes of hoarseness: (Table 2)(fig2)The commonest cause of hoarseness in the present study was found to be acute Laryngitis (26.66%) followed by vocal cord polyp(22.5%) and vocal cord nodule(18.33%).3- Distribution of causes of hoarseness

acrosssex(table3 figure3)Analysis of sex data using chi-square test revealed that acute laryngitis vocal cord , nodule and chronic non specific laryngitis affects significantly males more than females (P-value < 0.001). In contrast to vocal cord polyp, which significantly affects females more than males (P - value < 0.0001).

Regarding the remaining causes of hoarseness, there is no significant difference in incidence between males and females Table (3

)distribution of cases of hoarseness across sex Wile figure 3 showing that the incidence of hoarseness is more common between male(55%)

hoarseness presenting in less than 2 weeks and chronic hoarseness presenting for more than 2 weeks.as shown in figure 1,cases with chronic hoarseness(65.83)

gender than female gender (45%) 4. Distribution of causes of hoarseness across age groups: (Table

4)Patients were categorized into seven age groups (0-10, 11-20, 21-30, 31- 40, 41-50,51-60,and 61-70).Analysis of data shows that the mean age of presentation,in the present study is 40 years for the total sample,About 55.5% of patients were younger than 40 years and,44.5% were older than 40 years. For the total

sample,Regarding acute Laryngitis of cases fall in two age groups (21-30 and 31-40 years),In vocal cord polyp, maximum of cases occur in age group (31-40years) .5- Patients'

residence : Whether urban or rural is shown in figure and table (5) which clearly demonstrates that laryngeal pathologies which cause hoarseness affect predominantly those who live in urban59.16% rather than rural area 40.83%),,6-

Prevalence across occupational groups, Table (6)and fig(6) shows that maximum incidence of hoarseness where between

teachers(20%)and housewives(16.66%) and less between engineer and military men
 7- Relationship between smoking and vocal abuse and the development of laryngeal pathologies causing hoarseness of voice : (Table 7)A significant relationship has been found

between smoking ,vocal abuse and development of laryngeal polyp, acute laryngitis vocal cord nodule and chronic non specific laryngitis (p-value < 0.0001) in contrast to the remaining conditions which show no significant relationship.

table 1 acute and chronic hoarseness

Type of hoarseness	Number of patients	Percentage
Acute hoarseness	41	34.16%
Chronic hoarseness	79	65.84%





table 2 causes of hoarseness

Cause	A.M.Bahaalddin, A.H.Jamal disorders in Azadi teaching hospital	No	%
Acute laryngitis		32	26.66
vocal cord polyp		27	22.5
vocal cord nodule		22	18.33
chronic non specific laryngitis		18	15
psychogenic dysphonia		7	5.83
vocal cord strain		3	2.5
blunt trauma to the neck		3	2.5
vocal cord palsy		2	1.66
Reinke's oedema		2	1.66
laryngeal papilloma		2	1.66
Intubation injury		1	0.83
laryngeal TB		1	0.83
Total		120	100%



Cause	Males	P-value	females	P value
Acute laryngitis	20	0.0001	12	0.001
vocal cord polyp	12	0.0001	15	0.0001
vocal cord nodule	13	0.0001	9	0.003
chronic non specific laryngitis	11	0.001	7	0.05
psychogenic dysphonia	3	0.006	4	0.0022
vocal cord strain	1	0.2	2	0.09
blunt trauma to the neck	2	0.09	1	0.2
vocal cord palsy	0	-----	2	0.09
Reinke's oedema	1	0.29	1	0.29
laryngeal papilloma	1	0.29	1	0.29
Intubation injury	0	-----	1	0.29
laryngeal TB	0	-----	1	0.29
Total	64		56	

table 3 the differences between male and female in the incidence of hoarseness





table 4 distribution between age groups

Cause	0-10	11-20	21-30	31-40	41-50	51-60	61-70	total
Acute laryngitis	0	0	0	9	7	2	3	21
vocal cord polyp	0	0	10	5	2	2	2	22
chronic non specific laryngitis	0	0	1	3	9	2	3	18
Psychogenic dysphonia	0	0	3	2	1	1	0	7
vocal cord strain	0	0	2	1	0	0	0	3
blunt trauma to the neck	0	0	1	1	1	0	0	3
vocal cord palsy	0	0	0	1	1	0	0	2
Reinke's oedema	0	0	0	1	1	0	0	2
laryngeal papilloma	2	0	0	0	0	0	0	2
Intubation injury	0	0	0	1	0	0	0	1
laryngeal TB	0	0	0	1	0	0	0	1
Total	2	4	35	31	25	10	13	120

Fig 4 the distribution of hoarseness between age groups



Fig(5) the proportion of patients residence





Fig (6) prevalence of hoarseness across occupation



Occupation	No. of patients	%
Teachers	24	20%
House wives	20	16.66%
Students	12	10%
Workers	10	8.33%
Medical staff	9	7.5%
Driver	9	7.5%
Unemployed	9	7.5%
Retired	8	6.66%
Farmers	7	5.8%
Engineer	6	5%
Military man	6	5%
Total	120	100%

table 6 hoarseness and occupations

Cause	No. of patients	No. of smokers	p-value	No. of vocal abuse	P-value
Acute laryngitis	32	11	0.0001	8	0.0001
vocal cord polyp	27	15	0.0001	10	0.0001
vocal cord nodule	22	6	0.0001	18	0.0001
chronic non specific laryngitis	18	11	0.0001	2	0.0031
psychogenic dysphonia	7	2	0.0198	5	0.00091
vocal cord strain	3	0	-----	3	0.05
blunt trauma to the neck	3	1	0.2	0	-----
vocal cord palsy	2	0	-----	2	0.18
Reinke's oedema	2	2	0.18	2	0.18
laryngeal papilloma	2	0	-----	0	-----
Intubation injury	1	0	-----	0	-----
laryngeal TB	1	0	-----	0	-----
Total	120	47		50	

Table7 the relationship of smoking and voice abuse and

Discussion

1-Causes of hoarseness :In the present study, acute laryngitis was found to be the most common cause of hoarseness accounting for 26.66% of total sample. This agrees with AL-Samarrae study - (8) (18.75%) . But disagrees with Herrington - Hall et al ((9) who stated that vocal cord nodule is the commonest cause of hoarseness (21.5%) .But this value is mostly an underestimate of the real prevalence of this condition, probably because most patients complaining from acute laryngitis were treated by general practitioners without being referred to otolaryngologists, in addition to seasonal variation of

presentation which predominates in winter months in association with upper respiratory tract infectio. 2-Sex distribution: The male to female ratio for the total study was found to be 1:2 in contrast to Hemngton-Hall et al(9) , found that laryngeal pathologies were more common in females than males Laryngeal pathologies which were found to be more common in males than in females are Acute laryngitis, vocal cord nodule, chronic non specific laryngitis While pathologies which were more common in females than males are vocal cord polyp (1.2:1) psychogenic dysphonia(2:5:1),vocal cord strain(3:0)

and vocal cord palsy(2:0) 3- Age distribution The mean age at presentation in the present study was 40 years which is close to that of AL-Samarrae study (38 years) About 55.5% of patients were younger than 45 years and 44.5% of patients were older than this age in our study in comparison to Herrington-Hall et al (9) in which 57% of patients were over 45 years old and 43% younger than this age . But it is important to notice that certain diseases occur predominantly in certain age group followed by, housewives 16.66% ,and students10% Herrington-Hall et al(9) showed that retired persons are most frequently involved in voice disorders, followed by housewives , workers,and unemployed. Teachers and house wives may abuse their voice and this may be the predominant cause of hoarseness. patients in our study. With significant association found between vocal abuse and the development of , vocalcord polyp. acute laryngitis, vocal cord nodule, 6- Relationship with smoking and vocal abuse: In the present study,

group like recurrent respiratory papillomatosis occur mainly in pediatric age groups 4- Patients' residence In our study, residents in urban area constituted 59.16% against 40.83% who live in rural area. This most probably reflects the effect of environmental pollution and working conditions as predisposing factors to vocal pathology in urban areas more than in rural areas5- Distribution across occupations In the present study, teachers20% were the most frequently affected occupational smokers represented 47% of total sample similarly, AL-Samarrae(8) reported that 57.5% of his series were smokers.Smoking has been shown to be significantly related to Acute laryngitis,chronic Laryngitis vocal cord polyp and vocal cord nodule (P-value < 0.0001) which is agreed by many authors Vocal abusers constituted 50% of total number of of patients in our study. With significant association found between vocal abuse and the development of , acute laryngitis, vocal cord nodule, and vocal cord polyp (P-value < 0.0001).

Again these findings correlate positively with the opinions of Paul van der Broek and P.H. Damste(4)

7- Associated symptoms

The most frequently associated symptom in the present study was sore

throat which accounted for 63% of cases .While in AL-Samarrae(8) series cough was the commonest associated symptom and it accounts for only 46.2% of cases .

Conclusions

1. The mean age of patients in the present study was 40 years at presentation. 2. The peak age incidence is between 21-30 years at presentation. 3. The commonest cause of hoarseness was found to be acute laryngitis of Larynx which predominantly affects young males. 4. Males are affected more than females by Laryngeal pathologies which may present with hoarseness of voice. 5. The commonest occupational group presenting as hoarseness was teachers persons followed by housewives. 6. Gastroesophageal reflux should not be overlooked as a predisposing factor to chronic Laryngitis

Recommendations:

1. Further studies on a larger sample conducted over time may provide better understanding of the distribution of laryngeal pathologies among voice disordered population. 2. Education for vocal hygiene and preventive management is warranted for those patients at high risk of developing voice disorders, namely elderly, factory workers, housewives and professional voice users. 3. Avoidance of smoking and vocal abuse play an important role in the management of hoarseness. 4- Early management of hoarseness carry better prognosis and easier in the management.

The study is self-funded

There is no conflict of interest between the authors

References

- 1-Boehme G, Gross M. Phoniatrie und Paedaudiologie. Kursew fur HNO-Aerzte. Berlin: Universitatsklinikum Benjamin Franklin 2000.
 - 2-Mathieson L. Analysis and evaluation: instrumental and perceptual. In: Mathieson L (ed.). The voice and its disorders. London: Whurr Publishers, 2001: 403-66.
 - 3- Stemple J. Voice research: A clear view of voice production, 25 years of progress: the speaking voice. Journal of Voice. 1993; 7: 293-300.
 - 4-Paul Van de Broeke and Ph Damste. Understanding voice problems - a Physiological perspective for diagnosis and treatment, 2nd edn. Maryland: Williams & Wilkins, 1996.
 - 5-Laugvaite J. Adult voice screening. Journal of Speech and Hearing Disorders. 1972; 37: 147-679
 - 6- Russell A, Dates J, Greenwood K. Prevalence of voice problems in teachers. Journal of Voice. 1998; 12: 467-79.
 - 7- Herrington-Hall B, Lee L, Stemple J, Niemi K, McHone M. Description of laryngeal pathologies by age, sex, and occupation in a treatment-seeking sample. Journal of Speech and Hearing 1998:22-56
 - 8- Al-Samarrae, Furat et al. The causes and classification of voice disorders. In: Freeman M, Fawcus M (eds). Voice disorders and their management, 3rd ed. London: Whurr Publishers, 2000: 47-66
 - 9- Herrington Hall et al. Voice disorders. In: Tomblin JB, Morris HL, Spriestersbach DC (eds). Diagnosis in speech-language pathology. San Diego, CA: Singular Press, 1994: 247-306.
 - 10- Pannebacker M. Voice and recommendations for outcome studies. American Journal of Speech-Language Pathology. 1998; 7: 49-64.
 - 11- Pannebacker M. Treatment of vocal nodules: Options and outcomes. American Journal of Speech-Language Pathology. 1999; 8: 209-17.
 - 12- Ramig L, Verdolini K. Treatment efficacy: Voice disorders. Journal of Speech, Language and Hearing Research. 1998; 41: S101-111
 - 13- Bergner M, Bobbin RA, Carter WB, Gilson BS. The Sickness Impact Profile: Development and final revision of a health status measure. Medical Care. 1981; 19: 787-805.
 - 14- World Health Organization. International Classification of Functioning, Disability and Health (ICF). Geneva: WHO, 2020
- treatment techniques: A review

15- Schalen L, Eliasson I, Fex S, Kamme C, Schalen C. Acute laryngitis in adults: results of erythromycin treatment. *Acta Otolaryngology*. 1992; supplement: 37-55.
gastroesophageal reflux disease (GERD): A clinical investigation of 225

16- Koufman J. The Otolaryngologic manifestations of gastroesophageal reflux disease (GERD): an experimental investigation of the role of acid and pepsin in the development of laryngeal injury. *Laryngoscope*. 1991; 101: 1-

79.

17- Hanson D, Conley D, Jiang J, Hahrilis P. Role of esophageal pH recording in management of chronic laryngitis an overview. *Annals of Otolaryngology, Rhinology, and Laryngology*. 2000; 114:4-9.

18- Roy N, Bless DM. Personality traits and psychological factors in voice pathology: 5:44-88

